
Appendix 3: Total hip replacements with a proximal modular femoral neck

- Flow diagram of included studies
- Study details 1 (aspects of internal validity)
- Study details 2 (aspects of external validity)
- Study reported outcomes
- References



Study	Study design	Study details I (aspects of internal validity)									
		Allocation method and concealment	Blinding (surgeons/ patients/ assessors)	Prospective collection and evaluation purpose	Sample size needs clearly defined?	Primary Outcome specified? (yes/no)	Intention- to-treat analysis? (yes/no)	Consecutive patients series? (yes/no)	Group comparability assessed?	Controlling for con- founding?	Procedure period
Duwelius, 2013	Retrospective comparison of successive non-consecutive cohorts	Time period	No	Prospective inclusion and collection, retrospective purpose	No	No	No	Unclear	Yes	Multi- variable analysis for baseline differences	August 2005 – December 2009
Gerhardt, 2014	Retrospective matched group comparison	Stock/availa- bility at operating room	No	No	No	Radio- graphic measure- ments	Unclear	No	No	Matching (demo- graphics and surgery date)	March 2008 – July 2011
Gualdi, 2006	Retrospective comparison of successive consecutive cohorts	Time period	No	No	No	No	No	Yes	No	No	1999 – 2006
Sakai, 2010	Retrospective matched group comparison	Surgeon's preference	Assessors	No	No	Unclear	No	Unclear	Demographics and preoperative assessments	Matching (demo- graphics and disease severity)	January 1994 – July 1996

Study	No. Of replace- ments (no. of patients)	Mean age (SD, range)	SD, (%)		Mean length of FU (yrs; SD, range)	Follow- up com- pletion (%)	Prosthesis brands (new vs conventional)	Manufactur er	Site, surgeon	Hospital setting (designer/ university/ general)	Continent (country)
Duwelius, 2013	878 (878)	62 (10)	52.4	93.2	NA (2.0- NA)	53.7-58.0	M/L taper Kinective versus M/L taper	Zimmer	Single site single surgeon,	General	North America (US)
Gerhardt, 2014	170 (170)	65.0 (33- 88)	56.5	100	NA (NA)	NA	Alloclassic Zweymuller versus Profemur Z	Zimmer and Wright Medical	Single site, multiple surgeons	General	Europe (Nether- lands)
Gualdi, 2006	519 (NA)	NA	NA	NA	1 (NA)	NA	Profemur Z <i>versus</i> Centerplus	Wright Medical and Ceraver	Single site, multiple surgeons	General	Europe (Italy)
Sakai, 2010	163 (133)	53.8 (40- 73)	91.9	100% second- ary to develop- mental dysplasia	14.5 (13- 15)	92.9	Custom designed uncemented modular neck system (Anca system) versus uncemented anatomical metal cancellous femoral component	Cremascoli (Milan, Italy) and ESKA (Lubeck, Germany)	Single site, multiple surgeons	University	Asia (Japan)

		Harris Hip Score (mean, SD / range)		Flexion (mean, SD / range)		Abduction (mean, SD / range)		Dislocation (count, proportion)		Revision (count, proportion)	
		Modular	Standard	Modular	Standard	Modular	Standard	Modular	Standard	Modular	Standard
Moderate	Preoperative Postoperative	52 (13) 92 (10)	50 (13) 90 (15)	NA NA	NA NA	NA NA	NA NA	- 5 (0.8%)	- 5 (1 8%)	- 7 (1 4%)	- 3 (1.0%)
	rostoperative	32 (10)	JO (1J)	IVA	IVA	IVA	IVA	3 (0.070)	3 (1.070)	7 (1.470)	3 (1.070)
Low to	Preoperative	NA	NA	NA	NA	NA	NA	-	-	-	-
moderate	Postoperative	NA	NA	NA	NA	NA	NA	4 (4.2%)	4 (4.2%)	2 (2.1%)	2 (2.1%)
Low	Preoperative	NA	NA	NA	NA	NA	NA	-	-	-	-
LOW	Postoperative	NA	NA	NA	NA	NA	NA	8 (2.2%)	6 (3.9%)	NA	NA
	Prophorativo	44.6	46.6	71	72	13	14.9				
Low to moderate	rieoperative	(18.4-0.5)	(16.1-5.7)	(10-110)	(15-120)	(0-30)	(0-35)	-	-	-	-
	Postoperative	98.6* (64-100)	93.9 (68-100)	96 (45-120)	91 (5-120)	32* (15-40)	28 (0-40)	0 (0%)	0 (0%)	0 (0%)	1 (1.4%)
1	Low to noderate Low	Low to Preoperative Postoperative Postoperative Postoperative Postoperative Postoperative Postoperative Postoperative Preoperative Preo	Low to Preoperative NA Low Preoperative NA Preoperative NA NA Preoperative NA	Low to Preoperative NA	Low to Preoperative NA	Postoperative	Postoperative	Postoperative	Postoperative	Postoperative	Postoperative Postoperativ

NA = not available (not applicable or not provided), * significant difference

- Duwelius PJ, Burkhart B, Carnahan C, Branam G, Ko LM, Wu Y, et al. Modular versus nonmodular neck femoral implants in primary total hip arthroplasty: which is better? Clin Orthop Relat Res 2014 Apr;472(4):1240-5.
- Gerhardt DM, Bisseling P, de VE, van Susante JL. Modular Necks in Primary Hip Arthroplasty Without Anatomical Deformity: No Clear Benefit on Restoration of Hip Geometry and Dislocation Rate. An Exploratory Study. J Arthroplasty 2014 Feb 10.
- Gualdi A, Masnaghetti P, Borroni G, et al. Fixation and geometry in total hip prosthesis: Is compromise necessary or is optimisation possible?. [Italian]. Minerva Ortopedica e Traumatologica 57 (4):289-292, 2006.
- Sakai T, Ohzono K, Nishii T, Miki H, Takao M, Sugano N. A modular femoral neck and head system works well in cementless total hip replacement for patients with developmental dysplasia of the hip. J Bone Joint Surg Br. 2010 Jun;92(6):770-6.